

REFERENCES

- Barko, J. 1988. Interactions between macrophyte growth and sediment nutrient availability. U.S. Army Corps of Engineers Waterways Experiment Station. 28 pp.
- Borman, S. 1991. The distribution and density of aquatic plants in Lake Wissota, Chippewa County, Wisconsin. Wisconsin Department of Natural Resources. Eau Claire, Wisconsin.
- Chippewa River Settlement Team. 2001. Lower Chippewa River Settlement Agreement. FERC #1982; 2567; 2670; 2639; 2491; 2440
- Cooke, D. 2007. Lake level drawdown as a macrophyte control technique. *J. Amer. Water Res. Assn.* 16(2):317-322
- Delong, M. D. and N. D. Mundahl. 1995. Effects of late-winter drawdown on benthic invertebrate community structure in the littoral zone of Lake Wissota, Wisconsin. Wisconsin Department of Natural Resources. Eau Claire, Wisconsin.
- Duarte, C. M. & J. Kalff. 1986. Littoral slope as a predictor of the maximum biomass of submerged macrophyte communities. *Limnology Oceanographic.* 31(5): 1072-1080
- Dunst, R. 1982. Sediment problems and lake restoration in Wisconsin. *Environment International.* 7: 82-92
- Engel, S. 1985. Aquatic community interactions of submerged macrophytes. Wisconsin Department of Natural Resources Tech. Bulletin No. 156. 79pp.
- Engel, S. 1990. Ecosystem response to growth and control of submerged macrophytes: A literature review. Wisconsin Department of Natural Resources Tech. Bulletin No. 170. 20 pp.
- Haxton, T. J. and C. S. Findlay. 2008. Meta-analysis of the impacts of water management on aquatic communities. *Canadian Journal of Fisheries and Aquatic Sciences.* 65: 437-447.
- Howard, R and C. Wells. 2009. Plant community establishment following drawdown of a reservoir in southern Arkansas, USA. *Wetlands Ecol. Manag.*
- Heuschele, Deborah Jo. 2006. A comparison of the distribution and density of aquatic plants in Lake Wissota, Chippewa County, Wisconsin between 1989 and 2005. Beaver Creek Reserve Citizen Science Center.

- Kaster, J. L. and G. Z. Jacobi. 1978. Benthic macroinvertebrates of a fluctuating reservoir. *Freshw. Biol.* 8: 283-290.
- Konkel, D. 1998. Impacts of winter drawdowns on the aquatic plant community in Lake Wisconsin. Wisconsin Department of Natural Resources. Eau Claire, Wisconsin.
- Jessen, R. & R. Lound. 1962. An evaluation of a survey technique for submerged aquatic plants. Minnesota Department of Conservation Game Investigation Rep. No. 6. 10 pp.
- Leira, M. 2008. Effects of water level fluctuation on lakes: an annotated bibliography. 613 (1):171-184.
- Lombardo, P. & P.G. Cooke. 2003. Ceratophyllum demersum - phosphorus interactions in nutrient enriched aquaria. *Hydrobiologia.* 497(1-3): 79-90
- Moody, ML and DH Les. 2007. Geographic distribution and genotypic composition of invasive hybrid watermilfoil (*Myriophyllum spicatum* X *M. Sibiricum*) populations in North America.
- Nichols, S. 1999. Floristic quality assessment of Wisconsin lake plant communities with example applications. *Journal of Lake and Reservoir Management.* 15(2): 133-141
- Nichols, S., S. Weber, & B. Shaw. 2000. A proposed aquatic plant community biotic index for Wisconsin lakes. *Environmental Management.* 26(5): 491-502
- Nichols, S. 2008. The use of overwinter drawdown for aquatic vegetation management. *J. Amer. Water Res. Assn.* 11(6): 1137-1148.
- Richardson, S. M., J. M. Hanson and A. Locke. 2002. Effects of impoundment and water-level fluctuations on macrophyte and macroinvertebrate communities of a dammed tidal river. *Aquatic Ecology.* 36: 493-510.
- Wetzel, R.G. 2001. *Limnology: Lake and River Ecosystems.* Third Edition. Academic Press.
- White, M., M. Xenopoulos, K. Hogsden, R. Metcalf and P. Dillon. Natural lake level fluctuation and associated concordance with water quality and aquatic communities within small lakes of the Laurentian Great Lakes region. *Hydrobiologia.* 613(1): 21-21.